



## R&D Systems ELISA Reference Guide & Catalog

**Featuring** ✓ Published References ✓ Insight from R&D Systems ELISA Experts  
✓ Assay Principles ✓ Troubleshooting Guide



To request your copy, click here or visit our website at [www.RnDSystems.com/go/request](http://www.RnDSystems.com/go/request)



### Sequential Binding of $\alpha_v\beta_3$ and ICAM-1 Determines Fibrin-Mediated Melanoma Capture and Stable Adhesion to CD11b/CD18 on Neutrophils

This information is current as of January 13, 2011

Pu Zhang, Tugba Ozdemir, Chin-Ying Chung, Gavin P. Robertson and Cheng Dong

*J Immunol* 2011;186:242-254; Prepublished online 6 December 2010;  
doi:10.4049/jimmunol.1000494  
<http://www.jimmunol.org/content/186/1/242>

- 
- |                      |  |
|----------------------|--|
| <b>References</b>    | This article <b>cites 55 articles</b> , 34 of which can be accessed free at: <a href="http://www.jimmunol.org/content/186/1/242.full.html#ref-list-1">http://www.jimmunol.org/content/186/1/242.full.html#ref-list-1</a> |
| <b>Subscriptions</b> | Information about subscribing to <i>The Journal of Immunology</i> is online at <a href="http://www.jimmunol.org/subscriptions">http://www.jimmunol.org/subscriptions</a>   |
| <b>Permissions</b>   | Submit copyright permission requests at <a href="http://www.aai.org/ji/copyright.html">http://www.aai.org/ji/copyright.html</a>  |
| <b>Email Alerts</b>  | Receive free email-alerts when new articles cite this article. Sign up at <a href="http://www.jimmunol.org/etoc/subscriptions.shtml/">http://www.jimmunol.org/etoc/subscriptions.shtml/</a>                              |

---

*The Journal of Immunology* is published twice each month by The American Association of Immunologists, Inc., 9650 Rockville Pike, Bethesda, MD 20814-3994. Copyright ©2011 by The American Association of Immunologists, Inc. All rights reserved. Print ISSN: 0022-1767 Online ISSN: 1550-6606.



# Sequential Binding of $\alpha_v\beta_3$ and ICAM-1 Determines Fibrin-Mediated Melanoma Capture and Stable Adhesion to CD11b/CD18 on Neutrophils

Pu Zhang,\* Tugba Ozdemir,\* Chin-Ying Chung,<sup>†</sup> Gavin P. Robertson,<sup>†</sup> and Cheng Dong\*

**Fibrin (Fn) deposition defines several type 1 immune responses, including delayed-type hypersensitivity and autoimmunity in which polymorphonuclear leukocytes (PMNs) are involved. Fn monomer and fibrinogen are multivalent ligands for a variety of cell receptors during cell adhesion. These cell receptors provide critical linkage among thrombosis, inflammation, and cancer metastasis under venous flow conditions. However, the mechanisms of Fn-mediated interactions among immune cells and circulating tumor cells remain elusive. By using a cone-plate viscometer shear assay and dual-color flow cytometry, we demonstrated that soluble fibrinogen and Fn had different abilities to enhance heterotypic aggregation between PMNs and Lu1205 melanoma cells in a shear flow, regulated by thrombin levels. In addition, the involvement of integrin  $\alpha_v\beta_3$ , ICAM-1, and CD11b/CD18 (Mac-1) in fibrin(ogen)-mediated melanoma-PMN aggregations was explored. Kinetic studies provided evidence that ICAM-1 mediated initial capture of melanoma cells by PMNs, whereas  $\alpha_v\beta_3$  played a role in sustained adhesion of the two cell types at a shear rate of  $62.5\text{ s}^{-1}$ . Quantitative analysis of the melanoma-PMN interactions conducted by a parallel-plate flow chamber assay further revealed that at a shear rate of  $20\text{ s}^{-1}$ ,  $\alpha_v\beta_3$  had enough contact time to form bonds with Mac-1 via Fn, which could not otherwise occur at a shear rate higher than  $62.5\text{ s}^{-1}$ . Our studies have captured a novel finding that leukocytes could be recruited to tumor cells via thrombin-mediated Fn formation within a tumor microenvironment, and  $\alpha_v\beta_3$  and ICAM-1 may participate in multistep fibrin(ogen)-mediated melanoma cell adhesion within the circulation. *The Journal of Immunology*, 2011, 186: 242–254.**

**M**elanoma cancer metastasis is a highly regulated process. Circulation-mediated metastasis requires lodging of cells to distinct sites within vasculature where melanoma cells extensively interact with extracellular environment including platelets, leukocytes, and plasma proteins. In face of fluid shear forces, melanoma cells need to express shear-resistant receptors to adhere to endothelial cells (ECs) of the vascular wall. Unlike leukocyte-endothelial cell binding, adhesion between melanoma cells and endothelial cells does not occur via direct receptor-ligand binding, because most of melanoma cells do not express  $\beta_2$  integrins or Sialyl Lewis X (sLe<sup>x</sup>) at the levels capable of facilitating the binding to the endothelium (1). Previous studies have reported that platelets and polymorphonuclear leukocytes (PMNs) could facilitate hematogenous dissemination of mel-

anoma cells by seeding them to the endothelial wall (1–3). However, it is still not well understood how plasma proteins, especially fibrinogen (Fg) or fibrin (Fn) expressed within a tumor microenvironment, may regulate tumor cell adhesion.

A clear link between hemostasis and tumor metastasis has been discovered by both in vivo and in vitro studies. For example, cancer patients were often shown to have abnormalities of blood coagulations, with elevated levels of Fg and fibrinopeptide A, which is a byproduct of Fn formation (2). Melanoma cells secrete tissue factors, which are the precursors of coagulation cascade leading to Fn production that promotes metastasis by mediating prolonged adhesion of melanoma cells to endothelial cells (4). Effects of Fn monomers on binding of platelets to melanoma cells under flow conditions were examined by several investigators (2, 3, 5). In these studies, Fn was shown to enhance platelet and melanoma cell aggregation by bridging integrin  $\alpha_{IIb}\beta_3$  on platelets to either integrin  $\alpha_v\beta_3$  (3) or ICAM-1 (2, 5) on melanoma cells. A similar mechanism was seen in Fg-enhanced leukocyte-endothelial cell adhesion through binding of Fg to ICAM-1 on endothelial cell (6). Importantly, shown in an animal model of experimental metastasis compared with wild-type controls, depletion of Fg reduced sustained adhesion of tumor cells, whereas treating tumor cells with soluble Fn enhanced lung seeding within the microcirculation (2, 7). Recently, platelet-carcinoma heterotypic aggregation in a suspension under shear conditions was also shown to be interfered with by soluble Fn, which bound to CD44 on carcinoma cells and diminished CD44-P-selectin interactions (8). Therefore, the enhancement of tumor metastasis by Fg and Fn may be viewed from heterotypic cell-cell adhesion influenced by both mechanical shear and kinetic binding mechanisms.

The fluid shear in circulation that facilitates cell-cell collisions can be translated to a tensile shear force in breaking cellular aggregates (9). Cell-cell adhesion is additionally affected by the

\*Department of Bioengineering, The Pennsylvania State University, University Park, PA 16802; and <sup>†</sup>Department of Pharmacology, College of Medicine, The Pennsylvania State University, Hershey, PA 17003

Received for publication February 16, 2010. Accepted for publication October 21, 2010.

This work was supported by National Institutes of Health Grant CA-127892, American Cancer Society Grant RSG-04-053-01-GMC, and the Foreman Foundation for Melanoma Research (to G.P.R.) and by National Institutes of Health Grant CA-125707 and National Science Foundation Grant CBET-0729091 (to C.D.).

Address correspondence and reprint requests to Dr. Cheng Dong, The Pennsylvania State University, 233 Hallowell Building, University Park, PA 16802. E-mail address: cxd23@psu.edu

Abbreviations used in this paper: CD44s, standard form of CD44; DPBS, Dulbecco's PBS; EC, endothelial cell; EI, fibroblast L cell that had been transfected to express human E-selectin and ICAM-1; Fg, fibrinogen; fMLF, N-formyl-methionyl-leucyl-phenylalanine; Fn, fibrin; GPRP, Gly-Pro-Arg-Pro amide; PMN, polymorphonuclear leukocyte; siRNA, small interfering RNA; sLe<sup>x</sup>, Sialyl Lewis X; TP, single tumor cell with one polymorphonuclear leukocyte; TP2, single tumor cell with two polymorphonuclear leukocytes; TP3+, single tumor cell with more than three polymorphonuclear leukocytes; TRITC, tetramethylrhodamine isothiocyanate isomer R.

Copyright © 2010 by The American Association of Immunologists, Inc. 0022-1767/10/\$16.00

intrinsic binding properties between receptors and ligands on respective cell types. Therefore, without a proper affinity between receptors and ligands, cell–cell adhesion cannot occur in an ordered fashion. Most melanoma cells have been shown to lack necessary selectins and integrins that are responsible for the tethering/retaining of tumor cells to the endothelial wall within the circulation. Instead, melanoma cells do express high levels of ICAM-1 that can bind to  $\beta_2$  integrins such as LFA-1 (CD11a/CD18) and Mac-1. Recent studies have shown that ICAM-1–expressing melanoma cell could effectively adhere to PMNs in a very cooperative and sequential process, consisting of LFA-1–mediated initial capture and Mac-1–dependent firm adhesion (10).

In the presence of Fg or Fn, adhesion of melanoma cells to PMNs via fibrin(ogen) might follow a similar process mediated by bridging ICAM-1 (on melanoma) to fibrin(ogen) to  $\beta_2$  integrins (on PMNs), which is the main focus of this paper. Besides ICAM-1, melanoma cells express  $\alpha_v\beta_3$ , which can bind to Fg and Fn both under static and flow conditions (11–13). In melanoma cells, expression of  $\alpha_v\beta_3$  is associated with metastatic phenotypes. Fg contains three potential  $\alpha_v\beta_3$  binding sites at A $\alpha$ 95–98 (RGDF), A $\alpha$ 572–575 (RGDS), and  $\gamma$ C400–411 (12, 13). Many cellular interactions with Fg and Fn occur via binding to one or two of these recognition sequences. Upon thrombin cleavage of fibrinopeptides A, Fg may expose more cryptic RGD sites that mediate stronger  $\alpha_v\beta_3$  binding. The major ICAM-1 recognition site is located in  $\gamma$ 117–133 of Fg (14). The leukocyte integrin Mac-1 is a high-affinity receptor for Fg on stimulated monocytes and PMNs, which has been implicated in many inflammatory responses. Mac-1 interaction site is localized within Fg D domain at a site corresponding to the  $\gamma$ -chain 190–202 and 377–395 (15).

The binding kinetics of melanoma cells and PMNs under hydrodynamic conditions has been recently investigated in the absence of fibrin(ogen) by using a cone-plate viscometer assay (16). The cell collision frequency, duration of cell–cell contact, and shear stresses acting on the cell could also be theoretically determined (9, 17). Because PMNs were shown to be involved in tumor cell extravasation and Fn levels were also shown an increase in cancer patients affected by plasma-contained thrombin (7, 18), we hypothesized in the current study that thrombin-regulated soluble Fn formation might mediate specific processes of melanoma–PMN binding by bridging the two cell types. In this paper, we have examined the binding of melanoma cells to Fn or Fg and its subsequent effects on melanoma–PMN adhesion. In addition, we have provided a quantitative comparison of relative roles in ICAM-1 and  $\alpha_v\beta_3$ –mediated melanoma–PMN interactions under various hydrodynamic shear conditions. We have found for the first time, to our best knowledge, that Fg and Fn-enhanced melanoma–PMN binding lies in a potential mechanism of ICAM-1–mediated initial capture followed by  $\alpha_v\beta_3$ –enhanced firm retention of melanoma adhesion to PMNs via fibrin(ogen). This enhancement of PMN–melanoma binding resulted in more melanoma being brought into close proximity of ECs, thereby representing a prelude of melanoma adhesion-led invasion and metastasis.

## Materials and Methods

### Abs and reagents

Mouse IgG anti-human mAbs against  $\alpha_v\beta_3$  (anti-CD51/61, clone 23C6) and ICAM-1 (clone BBIG-11) were purchased from R&D Systems (Minneapolis, MN). Mouse anti-human mAbs against LFA-1 (anti-CD11a) and Mac-1 (anti-CD11b) were purchased from Invitrogen (Carlsbad, CA). *N*-formyl-methionyl-leucyl-phenylalanine (fMLF), Fg (fraction I, type I from human plasma), Gly-Pro-Arg-Pro amide (GPRP), tetramethylrhodamine isothiocyanate isomer R (TRITC), chondroitinase AC II (*Arthrobacter aurescens*), and BSA were purchased from Sigma-Aldrich (St. Louis, MO). Fibronectin was obtained from VWR (West Chester, PA). Human serum

albumin was purchased from Calbiochem (La Jolla, CA). Thrombin bovine (269,300 U/g) was purchased from MP Biomedicals (Solon, OH). LDS-751 was purchased from Invitrogen.

### Cell culture and preparation


A375m, Lu1205, and WM35 (provided by Dr. Meenhard Herlyn, Wistar Institute, Philadelphia, PA) melanoma cells were grown in DMEM Nutrient Mixture F12 and RPMI 1640 (Life Technologies, Carlsbad, CA), respectively, supplemented with 10% FBS (BioSource International, Carlsbad, CA). Prior to each experiment, Lu1205 cells were detached with 0.05% trypsin/EDTA (Invitrogen) and washed twice with fresh medium. The cells were then suspended in fresh media and allowed to recover for 1 h while being rocked at a rate of 8 rpm at 37°C. [In this way, ICAM-1 expression was not affected, whereas integrin  $\alpha_v\beta_3$  could be regenerated after 1 h recovery (Fig. 1A, 1B)]. In some receptor-blocking experiments, Lu1205 cells were pretreated with respective functional blocking Ab (e.g., anti- $\alpha_v\beta_3$  [5  $\mu$ g/ml], anti-hICAM-1 [5  $\mu$ g/ml]) for 30 min at 37°C, preonset of assays. In selective experiments, to prevent CD44 binding to Fn, specific glycosaminoglycans from CD44 were removed by pretreating melanoma cells with 0.1 U/ml chondroitinase AC II [which catalyzed the cleavage of *N*-acetylhexosaminide linkage in chondroitin sulfate and has been shown to significantly suppress binding of CD44-expressing cells to Fn (19)] for 1 h at 37°C.

PMN preparation was previously described (1). Following The Pennsylvania State University Institutional Review Board-approved protocols (number 19311), we collected fresh human blood from healthy adults by venipuncture. PMNs were isolated using a Ficoll-Hypaque (Histopaque, Sigma-Aldrich) density gradient as described by the manufacturer and kept at 4°C in Dulbecco's PBS (DPBS) containing 0.1% human serum albumin for up to 4 h before an experiment. In selected experiments, PMNs were functionally blocked with anti-LFA-1 (5  $\mu$ g/ml) and anti-Mac-1 (5  $\mu$ g/ml).

Fibroblast L cells that had been transfected to express human E-selectin and ICAM-1 (EI cells; kindly provided by Dr. Scott Simon, University of California Davis, Davis, CA) were maintained in culture as described elsewhere (1, 20). EI cells express ICAM-1 at a level comparable with IL-1 $\beta$ –stimulated HUVECs (21) and were used as a stable ICAM-1–expressing endothelial monolayer for some of the cell adhesion studies.

### Preparation of soluble fibrin solution

Soluble Fn was made freshly in ion-free DPBS prior to each experiment. To produce soluble Fn monomers and prevent coagulation, thrombin cleavage of Fg was initiated in the presence of 4 mM GPRP-NH<sub>2</sub> following



**FIGURE 1.** The effect of trypsinization on receptor expression. Lu1205 cells were detached from culture dishes by 0.05% trypsin-EDTA. Cells were incubated for 60 min in medium before being treated with trypsin for 5 min. ICAM-1 (A) and  $\alpha_v\beta_3$  (B) expressions on cells at 0 min and 60 min after being detached as well as 5 min posttrypsinization were detected by flow cytometry. There were no significant statistical differences between groups.

a published protocol (19, 22). To probe the kinetics of Fn production, thrombin concentrations were varied while keeping a chosen dose from soluble Fg and GPRP. For example, to make 1 ml Fn solution, 120  $\mu$ l Fg (25 mg/ml) and 168  $\mu$ l GPRP (24 mM) were mixed with either 0  $\mu$ l, 2.7  $\mu$ l, 5.5  $\mu$ l, or 207  $\mu$ l thrombin (10 U/ml, 269,300 U/g) immediately preincubation at 37°C for 10 min. Then, a 2-fold concentrated Fn stock solution was subsequently mixed with cell suspension (containing tumor cells and/or PMNs) 1 s pre-experiment at a 1:1 ratio to reach a desired Fg, thrombin, GPRP, and cell concentrations.

#### SDS-PAGE for fibrinogen-digested products

To characterize thrombin catalyzed digestion of Fg (3 mg/ml), the reactions were run at room temperature and terminated at selected intervals by adding 2 $\times$  SDS running buffer (0.2% bromophenol blue, 4% SDS, 100 mM Tris [pH 6.8], 20% glycerol) and 2-ME. The samples were analyzed by SDS-PAGE on 12% gels. Control was prepared by adding SDS and 2-ME to Fg before thrombin and GPRP. Gels were stained with Coomassie blue and imaged accordingly.

#### Small interfering RNA targeting ICAM-1 and integrin $\alpha_v\beta_3$

A total of 100 pmol duplexed Stealth small interfering RNA (siRNA; Invitrogen, Carlsbad, CA) was introduced into  $1.0 \times 10^6$  Lu1205 cells via nucleofection using an Amaxa Nucleofector using Solution R/program K-17. Transfection efficiency was listed in Table II with 80–90% cell viability. Following siRNA introduction, cells were replated in culture dishes. siRNA sequences were: scrambled: 5'-AAUUCUCCGAACGUGUCACGUGAG-A-3'; ICAM-1: 5'-UUAUAGAGGUACGUGUCAGGCCUG-3'; integrin  $\alpha_v$ : 5'-UUGAUGAGCUCAUAGACAUGGUGGA-3'; and integrin  $\beta_3$ : 5'-AUAAGCAUCAACAAUGASGUGGAGG-3'.

#### Binding of tumor cells or PMNs to immobilized fibrin or fibrinogen

Immobilized Fg surfaces were generated by absorbing Fg (von Willebrand factor-, plasminogen-, and fibronectin-free) solution at 2.5 mg/ml in DPBS containing 1.5 mM  $\text{Ca}^{2+}$  onto the substrate of a 35-mm polystyrene petri dish (BD Biosciences, Franklin Lakes, NJ) overnight at 4°C. Similarly, immobilized Fn surfaces were generated by incubating the above-mentioned Fg solution with thrombin (2 U/ml) for 2 h at 37°C (8). The non-specific binding sites were further blocked by incubating Fg- or Fn-coated surfaces with DPBS-1% BSA for 1 h at room temperature.

To assess circulating cells binding to immobilized fibrin(ogen), a parallel-plate flow assay was adapted following previously published protocols (1), in which suspension of either tumor cells or PMNs ( $5 \times 10^5$ /ml) in DMEM with 0.1% BSA was perfused over the Fn (or Fg)-coated surface for 5 min. Real-time flow digital videos were recorded under a 10 $\times$  objective (0.48 mm<sup>2</sup>) to a PC memory by Streampix III (Streampix III, Norpix, Montreal, Quebec, Canada). The ability of cell binding to immobilized fibrin(ogen) surfaces was quantified by the total number of firmly adhering cells (e.g., adhesion >30 s) during a 5-min period of perfusion under each condition.

#### Parallel-plate flow assay

Use of a parallel-plate flow assay allowed for direct observation of interactions between PMNs and melanoma cells and determination of kinetic parameters regulating tumor-PMN binding under flow conditions (23). In the current study, PMN binding to immobilized tumor cells was first assayed to characterize whether  $\beta_2$  integrins on PMNs bind to  $\alpha_v\beta_3$  or ICAM-1-expressing melanoma cells in the presence or absence of soluble fibrin(ogen). The dishes seeded with melanoma cell monolayer served as a substrate in a parallel-plate flow chamber. The field of view was 800  $\mu$ m long and 600  $\mu$ m wide. DMEM alone was first perfused into the chamber to allow the melanoma-cell monolayer to reach equilibrium. fMLF-stimulated PMNs were then allowed to presettle on the monolayer under a low shear rate of 10 s<sup>-1</sup> for 2 min before an onset of prescribed experimental shear rates (62.5 s<sup>-1</sup> or 200 s<sup>-1</sup>).

To simulate physiological conditions and assess melanoma adhesion to ECs assisted by PMN, suspension of fMLF-stimulated PMNs and melanoma cells at a ratio of 1:1 in the presence of Fn was perfused over an EI monolayer. To evaluate the capacity of PMNs to tether melanoma cells to ECs, melanoma adhesion efficiency was employed to quantify melanoma adhesion to the EI monolayer as a result of the collision with pretethered PMNs. If more than one melanoma cell adhered to a PMN, we counted it as multiple aggregates (e.g., two aggregates if two melanoma cells adhered to one PMN). Melanoma adhesion efficiency was expressed by the following ratio:

$$\frac{\text{Number of melanoma cells arrested on the monolayer}}{\text{Number of melanoma collisions to PMNs}}$$

The adhesion efficiency was then classified based on the durations of melanoma arrests (>1 s, >3 s, and >5 s).

#### Cone-plate viscometer assay

To study binding of melanoma cells to PMNs both in suspension and under shear conditions, the flow assay was adapted using a cone-plate viscometer (RotoVisco 1, Haake, Hewington, NH) as previously described (16, 17, 24). This viscometer consists of a stationary plate and a 1°-free rotating cone, capable of generating linear velocity fields with a constant gradient (i.e., shear rate). So when cell suspension was subjected to a shear field within the instrument, the fast moving cells near the rotating cone collide with slowly moving cells near the plate (17). Cell-cell collision generated one bond initially. Once one bond formed, more bonds might form or existing bonds might dissociate when under shear for a longer time. The tensile force determined the dissociation rate of a doublet. To later quantify heterotypic cell aggregation, isolated PMNs and melanoma cells were stained with LDS-751 (red) and TRITC (orange), respectively, for 10 min at 37°C. Thereafter, PMNs were mixed with melanoma cells in DPBS containing  $\text{Ca}^{2+}/\text{Mg}^{2+}$  in the presence of 1  $\mu$ M fMLF to reach a concentration of  $5 \times 10^5$  cells/ml for each cell type (1:1). The cell suspension was allowed to equilibrate at 37°C for 2 min. Fibrin(ogen) was added, respectively, into the cell suspension 1 s preinitiation of shear. Shear rates were varied from 62.5–200 s<sup>-1</sup> for preset duration of time, ranging from 30–300 s. After shear applications, samples were fixed immediately with 1% formaldehyde.

#### Quantification of heterotypic cell aggregation

An ability of TRITC-labeled melanoma cells to form aggregates with LDS-751-labeled PMNs was determined by using the dual-color flow cytometry based on the cellular compositions, which were characterized by cells' forward scatters, side scatters, and fluorescence profiles (Fig. 2) (16). Because the two cell types were labeled with different dyes, their aggregates would appear in the coordinates, which are integral multiples of singlet fluorescence values. A total of 5000 events were collected for each case. By applying this method, heterotypic aggregates comprised of a single tumor cell (T) with either one PMN (TP1), two PMNs (TP2), or more than three PMNs (TP3+) could be characterized. The extent of heterotypic aggregation was defined as the percentage of bound tumor cells to PMNs in total amount of cells: % tumor cell in heterotypic aggregates =  $\{(TP1) + (TP2) + (TP3+)\} / \{(T) + (TP1) + (TP2) + (TP3+)\}$ .


#### Statistical analysis

Data were obtained from at least three independent experiments and expressed by the means. Statistical significance of difference between means was determined by using Student *t* test or ANOVA. Turkey's test was used for post hoc analysis for ANOVA. Probability values of  $p < 0.05$  were chosen as statistical significance.

## Results

### Arrest of melanoma cells and PMNs on immobilized fibrin (ogen) under dynamic flow conditions

Melanoma cells express ICAM-1 and  $\alpha_v\beta_3$ , the receptors for fibrin (ogen), for which levels correlate with their respective metastatic potentials (Table I) (11, 25). The way to prepare Lu1205 did not affect the expression of these receptors (Fig. 1A, 1B). To analyze how melanoma cells or PMNs bind to fibrin(ogen) ligands under different flow conditions, a parallel-plate flow system was used where immobilized ligands were present on the substrate of the bottom plate. Perfusion of cell suspension over an immobilized Fn- or Fg-coated surface avoided some confounding factors and analysis for cell-cell and cell-soluble protein interactions in a three-dimensional flow field as shown in the following cone-plate assay (Fig. 2). When suspended in a binding buffer (DMEM + 0.1% BSA + HEPES), Lu1205 melanoma cells could bind to both immobilized Fn and Fg avidly at both low (62.5 s<sup>-1</sup>) and high (200 s<sup>-1</sup>) shear rates (Fig. 3A), although significantly more strongly to Fn than to Fg under low shear condition. Under both shear conditions tested,



**FIGURE 2.** Flow cytometric detection of time-dependent size growth of PMN-melanoma aggregates in the presence of Fn solution made from 0.053 U/ml thrombin, 1.5 mg/ml Fg, and 4 mM GPRP. *Top panels*, The TRITC-labeled Lu1205 population was resolved into singlet and aggregates composed of TP1, TP2, or TP3+ after 30 s (*left panels*), 180 s (*middle panels*), and 300 s (*right panels*) of shearing at shear rate of  $62.5 \text{ s}^{-1}$ . *Bottom panels*, Lu1205 cells were incubated with 1  $\mu\text{g/ml}$  anti-ICAM-1 mAb for 1 h before being labeled with TRITC-linked anti-mouse IgG. Similar to the *top panel*, these Lu1205 cells were subjected to shear with LDS751-labeled PMNs in the presence of Fn solution for 30 s (*left panels*), 180 s (*middle panels*), and 300 s (*right panels*) at a shear rate of  $62.5 \text{ s}^{-1}$ . Lu1205 singlet populations (*top panels*) over the time course were not those Lu1205 cells with low ICAM-1 expression; otherwise the singlet would have clustered to the left (*bottom panel*). They had the potential to bind to PMNs.

Lu1205 melanoma cells directly arrested on immobilized Fn or Fg surface without an apparent rolling step. To show the specificity of interactions, Lu1205 cells were also perfused over a BSA-coated surface (as a control in Fig. 3A), which resulted in minimal surface-bound arrests, suggesting melanoma cell binding to fibrin(ogen) shown in Fig. 3A was protein specific.

To further illustrate cell-surface receptors are responsible for binding, Lu1205 were perfused over immobilized Fn at shear rate of  $62.5 \text{ s}^{-1}$  in the presence of functional blocking mAbs, respectively, against ICAM-1,  $\alpha_v\beta_3$ , or IgG isotype control ( $5 \mu\text{g}/1 \times 10^6$  cells). As shown in Fig. 3B, the ability of Lu1205 cells to arrest on Fn under flow was only marginally affected by ICAM-1 blocking Ab, but much more effectively inhibited by anti- $\alpha_v\beta_3$  mAb, which demonstrates that  $\alpha_v\beta_3$  could be a more potent receptor than ICAM-1 on melanoma cells to support tumor cell adhesion to Fn and to make shear-resistant bonds.

PMNs were indicated to be capable of binding to either Fg- or Fn-coated surface under flow conditions (26, 27). To examine the effect of hydrodynamic shear force and chemoattractant stimulation on PMN binding to fibrin(ogen), nonstimulated or fMLF-stimulated ( $1 \mu\text{M}$ , 2 min) PMNs were perfused over an Fg- or Fn-coated surface in a parallel-plate flow chamber at a shear rate

of  $62.5 \text{ s}^{-1}$ . As shown in Fig. 3C, nonstimulated PMNs had a higher affinity for Fn than for Fg during the 5 min perfusion time, whereas they failed to bind to the BSA-coated surface (as a control; data not shown). In comparison, fMLF, a potent activator for the high-affinity state of  $\beta_2$  integrin, significantly enhanced PMN adhesion to both Fg and Fn (Fig. 3C). Like nonstimulated PMNs, activated PMNs bound to Fn with a higher affinity ( $220 \text{ PMN}/\text{mm}^2$ ) than to Fg ( $90 \text{ PMN}/\text{mm}^2$ ). Functional blocking Mac-1 Ab ( $5 \mu\text{g}/1 \times 10^6$  cells; for 30 min after being stimulated by fMLF) inhibited PMN binding to fibrin(ogen), which was in an agreement with previous studies (26). Taken together, results from Fig. 3C suggest that PMN adhesion to fibrin(ogen) is Mac-1 dependent.


#### *Tumor cells binding to PMNs is affected by thrombin-mediated Fn formation*

Using two-color flow cytometric assays, the heterotypic aggregates between TRITC-labeled tumor cells and LDS-751-labeled PMNs were quantified by their respectively gated characteristic fluorescence intensities. Some representative fluorescence profiles are shown in Fig. 2. It should be noticed that the melanoma singlet during the shear period represents those with normal expression of ICAM-1. They continue to be recruited to and dissociate from PMNs.

Table I. Flow cytometry analysis of ICAM-1 and  $\alpha_v\beta_3$  expressions on WM35, A375m, and Lu1205 cells

Receptor	Geometric Mean Fluorescence		
	WM35 (Low Metastatic)	A375m (Median Metastatic)	Lu1205 (Highly Metastatic)
Control IgG	$3.75 \pm 0.03$	$3.75 \pm 0.03$	$3.25 \pm 0.49$
ICAM-1	$33.76 \pm 0.25$	$106.76 \pm 10.2$	$165.5 \pm 12.8$
$\alpha_v\beta_3$	$26.22 \pm 0.45$	$16.58 \pm 0.36$	$45.5 \pm 1.16$

Values are geometric mean fluorescence intensities  $\pm$  SEM ( $n > 3$ ).



**FIGURE 3.** Characterization of Lu1205 and PMN surface receptors for binding to fibrin(ogen) under hydrodynamic shear conditions. *A*, Flow differentially affects the arrest of Lu1205 to immobilized Fg and Fn. Lu1205 cells were perfused over immobilized Fn or Fg in a parallel-plate flow chamber. The numbers of firm adhering cells ( $>30$  s) were quantified. Data represent means  $\pm$  SEM ( $n > 3$ ).  $*p < 0.05$  compared with respective Fg cases under different shear rates;  $^{\S}p < 0.05$  compared with control cases. Control cases quantified the numbers of Lu1205 adhesion to coated BSA (there were 0 cells adhering to BSA per  $\text{mm}^2$ ). *B*, Lu1205 cell adhesion to Fn requires  $\alpha_v\beta_3$  and ICAM-1. Data represent means  $\pm$  SEM ( $n > 3$ ).  $*p < 0.05$  compared with the isotype control case. *C*, Mac-1 mediates binding of PMNs to fibrin(ogen) under flow. Data represent means  $\pm$  SEM ( $n > 3$ ).  $*p < 0.05$ .

Binding rate of soluble proteins in solution to cell receptors follows an exponential-decay function of protein concentration (28). To approximately find the concentration-dependent Fg binding to melanoma cell and/or PMN receptors, the levels of melanoma-PMN aggregations mediated by soluble Fg with three arbitrary concentrations of 0.5, 1.5, and 3 mg/ml were compared (Fig. 4A). A total of 1.5 mg/ml Fg facilitated a significantly higher percentage of Lu1205-PMN heterotypic aggregation than 0.5 mg/ml Fg at 60 s. However, 3 mg/ml Fg did not further increase the extent of Lu1205-PMN aggregation compared with 1.5 mg/ml Fg concentration, suggesting a possible Fg saturation in mediating receptor-protein binding. Therefore, a 1.5 mg/ml concentration was chosen for probing Fg-mediated binding kinetics for the current study.

To understand potential mechanisms of Fn-mediated melanoma-PMN aggregation, Fn monomers were produced by reacting Fg, thrombin, and 4 mM GPRP. Conversion of Fg into Fn is initiated by thrombin-mediated cleavage in two NH<sub>2</sub>-terminal fibrinopeptides A and B (29). However, the concentrations of thrombin and

the reaction time would affect the kinetics of fibrinopeptide and Fn generation, and the amounts of generated Fn would have an impact on cell-cell interactions. To examine Fg-digested products, the components of Fg-thrombin reaction products were separated by SDS-PAGE to analyze the purity of Fn and mechanisms of thrombin catalysis. A total of 3 mg/ml Fn was cleaved by either 2 U/ml or 0.05 U/ml thrombin in the presence of GPRP for 10 min. The reactions were stopped by adding SDS and 2-ME. Proteins were resolved in 12% SDS-PAGE and stained with Coomassie blue (Fig. 4B). Fg  $\alpha$ -,  $\beta$ -, and  $\gamma$ -chains were present on the gel as separate bands, and their molecular weights matched well with proposed data (29). After thrombin cleavage, smaller m.w.  $\alpha$ - and  $\beta$ -chains were generated. In addition, the longer the time and the more thrombin, the more digested products were produced. Thrombin with a concentration of 2 U/ml could cleave all  $\alpha$ - and  $\beta$ -chains within 10 min.


To understand the kinetics of melanoma-PMN aggregation, the percentage of melanoma cells recruited by PMNs was plotted as a function of time. In the absence of soluble Fg (0 mg/ml), the number of aggregates increased with time and maximized at 60 s, and then rapidly decreased (Fig. 5A). This behavior was similar to WM9 melanoma-PMN binding found previously (16), which might be partially due to a rapid decrease in the cell concentration, with time leading to a decrease in cell collision frequency, or due to time-dependent reduction of cell receptor affinity (30).

Addition of Fg significantly elevated the percentage of heterotypic aggregation at shear rate of  $62.5 \text{ s}^{-1}$  (Fig. 5A). The percentage of Lu1205 in aggregation still peaked at 60 s and descended thereafter, similar to the case in the absence of Fg. Initial capture and subsequent stable adhesion between the two cell types increased proportionally in the presence of Fg (Fig. 5A) compared with cases without Fg. This implies that soluble Fg largely increased effective forward rate for receptor-ligand binding while being capable of maintaining the tumor cell-PMN aggregation.

Adding Fn solution (made from 0.025 U/ml thrombin and 1.5 mg/ml Fg) altered the biphasic trend of pure Fg-mediated tumor cell-PMN heterotypic aggregation at  $62.5 \text{ s}^{-1}$  (Fig. 5A). The presence of Fn allowed the binding to reach a steady state after the 60 s, increasing the percentage of aggregates from 48 to 57%, which sustained  $>5$  min. In comparison, Fn solution made by 0.053 U/ml thrombin boosted the initial melanoma-PMN aggregation to the same level (63%) of the Fg case over a period of 60 s, followed by a plateau without an apparent dissociation over at least 5 min. However, catalyzing Fg-Fn transformation using 0.025 U/ml or 0.053 U/ml thrombin might result in only a partial conversion, which could make an analysis of Fn-mediated binding mechanism more difficult. Therefore, we also used 2 U/ml thrombin to make apparently more completed Fn solution (8). Indeed, results showed that Fn solution made from 2 U/ml thrombin changed initial capture rate of Lu1205 cells compared with that made from 0.053 U/ml thrombin. A total of 70% Lu1205 cells were recruited to aggregates within 60 s. However, the level of prolonged aggregation was not changed. Collectively, these results suggest that thrombin concentration in a tumor microenvironment could be very important, and conversion of Fg to Fn exposed some cryptic recognition sites that may stabilize the formed aggregates or other forms of coagulation.

Percentage of aggregation decreased with an increase in shear rates (Fig. 5B). For example, during the first 60 s, 1000 cell-cell collisions resulted in 38 captures at a low shear rate of  $62.5 \text{ s}^{-1}$ , whereas 1000 collisions only resulted in 8 captures at a higher shear rate of  $200 \text{ s}^{-1}$ . Fg was able to upregulate the extent of adhesion of PMNs to Lu1205 melanoma cells under high shear of  $200 \text{ s}^{-1}$  to be comparable to that when Fg was absent under low

**FIGURE 4.** A, Effect of Fg concentration on PMN-Lu1205 binding. Lu1205 cells were subject to shear with PMNs in the presence of different concentrations of Fg in a cone-plate viscometer at a shear rate of  $62.5 \text{ s}^{-1}$ . Cell suspensions were fixed with formaldehyde at 60 s postonset of the experiments. Data represent means  $\pm$  SEM ( $n > 3$ ). \* $p < 0.05$ . B, SDS-PAGE for fibrinogen-digested products. Fg was cleaved by thrombin at different levels (0.05 U/ml [lane 3] and 2 U/ml [lane 4]) for 10 min. The reaction products were subjected to 12% polyacrylamide gel separation. Separated proteins were stained by Coomassie blue. Lane 1 is fibrinogen, lane 2 is control, and lane 5 is marker.



shear ( $62.5 \text{ s}^{-1}$ ) (Fig. 5A). According to the two-body collision theory, increasing shear rate may reduce capture efficiency due to an increase in fluid drag and tensile force in quickly rupturing transient bonds formation (9). Alternatively, increasing shear rate may also decrease the cell-cell encounter duration that limits the bond formation. Despite this transient feature of cell-cell aggregates at the shear rate of  $200 \text{ s}^{-1}$ , Fn was still able to slightly prolong the lifetime of aggregates. Fn solution made by 0.053 U/ml thrombin concentration shifted the peak binding from 60 s to 120 s at a higher shear rate of  $200 \text{ s}^{-1}$ . In general, Fn stabilized the transient process in tumor cell-PMN aggregation, given that it profoundly changed the time-dependent profile of fibrin(ogen)-mediated cell aggregation.

Compared with Lu1205 cells, WM35 cells, a low-metastatic melanoma cell line expressing lower amounts of receptors for fibrin(ogen) (Table I), failed to adhere efficiently to PMNs even in the presence of Fn at  $200 \text{ s}^{-1}$  (Fig. 5C), which correlated melanoma-PMN binding abilities with tumor metastatic potentials. Taken together, results from Fig. 5A-C suggested that the recruitment of PMNs to melanoma cells was dependent on thrombin-mediated Fn production in a tumor microenvironment with respect to tumor metastatic phenotypes and circulatory shear rates.


#### Relative roles of $\alpha_v\beta_3$ , ICAM-1, and Mac-1 in fibrin(ogen)-mediated binding

Lu1205 cells were incubated with a panel of  $\alpha_v\beta_3$  functional blocking Abs before being subjected to shear experiments in the cone-plate viscometer to determine the roles of  $\alpha_v\beta_3$  in heterotypic tumor cell-PMN aggregation.  $\alpha_v\beta_3$  blocking did not affect the rate of Lu1205 binding to PMNs in the absence of fibrin(ogen) (Fig. 6A), because  $\alpha_v\beta_3$  could not bind to  $\beta_2$  integrins on PMNs directly. Blocking  $\alpha_v\beta_3$  did not affect short-term binding between PMNs and Lu1205 melanoma cells in the presence of Fg (Fig. 6B). A total of 50% of Lu1205 cells were still able to bind to PMNs during the first 60 s, which could be due to the tethering via Fg between some unidentified receptors on the tumor cell and  $\beta_2$  integrins on PMNs, whereas the transient tethering quickly dissociated after 60 s, implying that Fg-dependent tethering did not sustain for longer time period during melanoma cell-PMN adhesion (Fig. 6B). Similarly, blocking  $\alpha_v\beta_3$  on melanoma cells had a marginal effect on initial Fn-mediated capture of Lu1205 cells

to PMNs, whereas the percentage of aggregates was significantly reduced compared with nonblocking cases after 120 s (Fig. 6C). Therefore, interaction of  $\alpha_v\beta_3$  with Fn-bound Mac-1 sustained Lu1205-PMN binding. Interestingly, Fn apparently prolonged the duration of  $\alpha_v\beta_3$ -independent transient tethering from 60 s compared with the Fg cases up to 120 s (Fig. 6A, 6C).

If  $\alpha_v\beta_3$  mediates the sustained adhesion, a question remains as what receptors on melanoma cells would play roles in tethering tumor cells to PMNs in the presence of soluble fibrin(ogen) under hydrodynamic shear. Fn  $\gamma$ -chain contains the ICAM-1 binding sites with a lower affinity than that of RGD sites for integrins (14). Therefore, we rationalized that ICAM-1 and Fn interactions might have a high on-rate and off-rate in bridging melanoma cells to PMNs, which would reduce the traveling speed of melanoma cells and allow a longer contact duration between the two cell types. To investigate the role of ICAM-1 on Fn-mediated melanoma cell and PMN binding, ICAM-1 was blocked by treating Lu1205 cells with functional blocking Abs anti-ICAM-1 before the shear experiments. Fig. 6A shows that blocking ICAM-1 eliminated native tumor cell-PMN aggregation, given that ICAM-1 was the predominant receptor on Lu1205 melanoma cells that facilitated PMN-melanoma binding (16). The 10% aggregates at 60 s might represent nonspecific binding. Addition of Fg increased the extent of ICAM-1-blocked Lu1205 in heterotypic binding to PMNs by 15%, and these aggregates lasted  $>5$  min under  $62.5 \text{ s}^{-1}$  shear condition, suggesting a strong ability of  $\alpha_v\beta_3$  to bind to Fg (Fig. 6B). Conversion of Fg to Fn further modified the  $\alpha_v\beta_3$ -dependent binding kinetics (Fig. 6C). Binding of  $\alpha_v\beta_3$  to Fn-bound Mac-1 slowly increased with time and reached a plateau at 120 s when 30% tumor cells were recruited to aggregates. These aggregates could sustain 5 min under shear without apparent disaggregation. However, blocking ICAM-1 reduced the rate of initial tethering of PMNs to Lu1205 cells in the presence of Fn by 60% (Fig. 6C, control versus anti-ICAM-1 case). This implies that the initial tethering (first 60 s binding postonset of shear) of Lu1205 cells to PMNs was due to ICAM-1 in terms of the high on-rate of ICAM-1-Fn binding.

LFA-1 has been shown to support the initial formation of melanoma-PMN aggregates (1). To distinguish LFA-1/ICAM-1-mediated initial tethering from potential Mac-1/Fn/ICAM-1-mediated events, we subsequently blocked LFA-1 on PMNs in ad-



**FIGURE 5.** Shear-induced adhesion of PMNs to highly-metastatic (Lu1205) and low-metastatic (WM35) melanoma cell lines in the presence of Fg or Fn. Fn solution was made from 1.5 mg/ml Fg and 4 mM GPRP with either 2 U/ml, 0.053 U/ml, 0.025 U/ml, or 0 U/ml thrombin. Control is PMN and melanoma cell binding in the absence of Fg or Fn. Percentage of Lu1205 cells bound to PMNs at shear rates of 62.5 s<sup>-1</sup> (A) and 200 s<sup>-1</sup> (B) are plotted as a function of time. C, Comparison of percentages of Lu1205 and WM35 bound to PMNs after 120 s of shearing at a shear rate of 200 s<sup>-1</sup>. \**p* < 0.05. Values are means ± SEM (*n* > 3).

dition to  $\alpha_v\beta_3$  blocking on Lu1205 cells before the shear experiments. It can be seen from Fig. 6A that blocking LFA-1 flattens the biphasic trend of Lu1205–PMN binding behavior. Although Fg slightly enhanced Mac-1–dependent binding (Fig. 6B) in a later time, Fn dramatically enhanced the initial binding (Fig. 6C), indicating that Fn added extra binding sites in tethering PMNs to Lu1205 cells. Taken together, LFA-1 cooperated with Fn-bound Mac-1 to enhance initial capture of tumor cells to PMNs in the presence of Fn.

Because there might be a potential concern about the Fc effect for Ab blocking, cells were subsequently transfected by siRNA to silence fibrin(ogen)-related receptors on Lu1205 cells,  $\alpha_v\beta_3$ , and ICAM-1 (knockout efficiencies were shown in Table II) before being subjected to shear experiments. As shown in Fig. 6D, in the presence of Fn,  $\alpha_v\beta_3$ -knockout Lu1205 cells remained being associated with PMNs for 120 s, and ICAM-1 supported Lu1205 aggregation with PMNs up to a peak level. However, after 120 s, disaggregation proceeded more rapidly than scramble and buffer cases. In contrast, rate of ICAM-1 knockout Lu1205 capture to

PMNs was significantly smaller than that of scramble and buffer cases, and  $\alpha_v\beta_3$ -mediated cell aggregation preceded very slowly and reached only 50% of the scramble and buffer cases at 5 min (Fig. 6E). These receptor knockdown results verified Ab blocking results, showing the relative roles of ICAM-1 and  $\alpha_v\beta_3$  in recruitment of Lu1205 cells to PMNs.

*Mac-1 is involved in fibrin binding, whereas LFA-1 and CD44 are not*


CD44 on carcinoma cells had an ability to bind to soluble Fn under flow conditions (8). Lu1205 melanoma cells express standard form of CD44 (CD44s; data not shown). Biochemical studies revealed that CD44s–fibrin(ogen) interaction was dependent on *N*-linked glycans that had a high affinity for Fn– $\beta$ -chain, which was distinct from Mac-1, ICAM-1, and  $\alpha_v\beta_3$  recognition sites on Fn (19, 31). To test whether other receptors, including CD44s, contribute to Fn-mediated PMN–Lu1205 heterotypic aggregation, ICAM-1 and  $\alpha_v\beta_3$  on Lu1205 cells were simultaneously blocked before shear experiments. Fig. 7A shows during 5 min shear,  $\alpha_v\beta_3$  supported 44% adhesion, ICAM-1 mediated 38% adhesion, whereas receptors other than these two supported 17% adhesion, indicating CD44s might not play as much roles as  $\alpha_v\beta_3$  and ICAM-1 in Fn-mediated PMN–Lu1205 melanoma cell heterotypic aggregation.

Pretreatment of PMNs with monoclonal anti-CD11b Ab inhibited the rate of their adhesion to Lu1205 cells via Fg by 50%, whereas the same Ab did not significantly affect the rate of their adhesion without Fg and slightly reduced longer term binding by 35% (Fig. 7B). This implied that Mac-1 participated in Fg- and Fn-mediated initial tether of PMNs to Lu1205, but only involved in sustained phase of native binding between PMN and Lu1205. Interestingly, at 60 s, when Mac-1 was blocked, Fg seemed to inhibit LFA-1 binding (anti-Mac-1 with 1.5 mg/ml Fg versus anti-Mac-1 without Fg case). This may result from the occupancy of free binding sites on ICAM-1 by Fg.

To further demonstrate the roles of ICAM-1 and  $\alpha_v\beta_3$  in Fn binding, while ruling out the role of CD44, adhesive phenotypes among several melanoma cells with different metastatic potential were compared. This included Lu1205 cells expressing ICAM-1 and  $\alpha_v\beta_3$ , A375 cells expressing high ICAM-1 (low  $\alpha_v\beta_3$ ) and WM35 cells expressing high  $\alpha_v\beta_3$  (low ICAM-1) (Table I). Before shear experiments, these cells were pretreated with chondroitinase AC II, which has been suggested to cleave binding motifs, like chondroitin sulfate and dermatan sulfate, on CD44 responsible for binding to Fn (19). Consistent with Ab blocking and siRNA knockout assays, the kinetics of WM35 binding mainly reflected that of  $\alpha_v\beta_3$ , whereas A375m binding was primarily contributed by ICAM-1 and reiterated the kinetics of ICAM-1 binding (Fig. 7C). Their adhesive phenotypes did not seem to be affected by enzymatic treatment targeting CD44.

*Fibrin(ogen) alters the mechanics of receptor-mediated melanoma cell adhesion to PMN and PMN-facilitated melanoma adhesion to ECs*

To further elucidate the relative roles of  $\alpha_v\beta_3$ , ICAM-1, and Mac-1 in fibrin(ogen)-mediated binding, direct video-microscopy observation of PMN–Lu1205 adhesion was obtained using a parallel-plate flow chamber assay in which a confluent Lu1205 cell monolayer served as a substrate of the flow chamber and PMNs were perfused over Lu1205 cells under two shear rates of 62.5 s<sup>-1</sup> and 20 s<sup>-1</sup>, respectively. The kinetics of PMN binding to immobilized Lu1205 cells assayed by a parallel-plate flow chamber was not identical to the binding kinetics characterized in cone-plate viscometer shear assays in which both PMNs and tumor cells were in suspension. This is because cone-plate assay data reflected a



**FIGURE 6.** Relative contribution of  $\alpha_v\beta_3$  and ICAM-1 to time-dependent adhesion of Lu1205 cells to PMNs in the absence of fibrin(ogen) (A), presence of fibrinogen (B), or fibrin (C–E) under  $62.5\text{ s}^{-1}$  shear rate. Lu1205 cells were pretreated with anti-ICAM-1,  $\alpha_v\beta_3$ , and/or CD11a mAb (A–C) or siRNA targeting ICAM-1 or  $\alpha_v\beta_3$  (D, E) before being sheared with PMNs. Fibrin was made from 0.053 U/ml thrombin, 1.5 mg/ml Fg, and 4 mM GPRP; fibrinogen stands for 1.5 mg/ml fibrinogen solution. Values are means  $\pm$  SEM from three independent experiments.

statistical snapshot of the aggregate formation at a chosen time point, whereas a parallel-plate assay provided a direct quantification in a real-time individual PMN–Lu1205 aggregation formation. For example, Fig. 8A clearly showed that Fg only enhanced the longer-period binding (i.e., sustained adhesion) between PMNs and Lu1205 melanoma cells (statistically significant for length of stop  $>3$  s), whereas Fn significantly enhanced both shorter- (i.e., initial tethered adhesion) and longer-period binding, which corroborated what we found in Fig. 8A when both cell types were in suspension under shear. In addition, in the presence of Fn, functional blocking of  $\alpha_v\beta_3$  on tumor cells significantly reduced firm adhesion (longer period  $>3$  s) of PMNs to Lu1205 cells, whereas blocking ICAM-1 on tumor cells only reduced initial capture ability (significant for time  $<3$  s) of Lu1205 cells under  $62.5\text{ s}^{-1}$  (Fig. 8B). This is again consistent with results from the cone-plate shear assay (Fig. 6C). Blocking ICAM-1 on Lu1205 cells did not significantly affect longer-period tumor cell binding to PMNs, suggesting it was the  $\alpha_v\beta_3$  that supported the firm adhesion in the presence of Fn.  $\alpha_v\beta_3$  on tumor cells was also shown to be sufficient in mediating optimal PMN–Lu1205 binding under very-low shear ( $20\text{ s}^{-1}$ ) which allows a longer contact time between PMN–Lu1205 (Fig. 8C). In contrast to Lu1205, non-metastatic WM35 melanoma cells failed to support appreciable PMN adhesion even in the presence of Fn (Fig. 8D), demonstrating the metastatic tumor specificity.

Table II. Transfection efficiency

Treatment	Geometric Mean Fluorescence	
	$\alpha_v\beta_3$	ICAM-1
Buffer	13	30.51
Scramble	12.14	40.32
siRNA	3.11	15
Knockout efficiency <sup>a</sup>	77%	62%


<sup>a</sup>Knockout efficiency is defined as the ratio of siRNA fluorescence intensity to scramble fluorescence intensity.

Fibrin(ogen) mediates melanoma–PMN aggregation, which may increase melanoma tethering to the endothelial wall and subsequently facilitate melanoma cell extravasation from the circulation (32). The mechanisms of bringing melanoma into close proximity to the endothelial cells by pretethered PMNs were studied in parallel plate flow assays with EI cells as a monolayer. In these experiments, it was observed that most of the melanoma cell binding to the EI under flow was enhanced by tethered PMNs. To normalize the resulting melanoma adhesion with respect to frequency of PMN–melanoma collision, a parameter, adhesion efficiency, was adopted (Materials and Methods, Fig. 8E). The time lengths of melanoma arrest on EI cells varied and were categorized to  $>1$  s,  $>3$  s, and  $>5$  s. Fn affected melanoma adhesion to EI cells via tethered PMN more than Fg did (Fig. 8E). The 1 s short-term melanoma–PMN tethering on EI cells was mediated by ICAM-1 on the melanoma (less by  $\alpha_v\beta_3$ );  $\alpha_v\beta_3$  maintained the bonds between PMN and melanoma, thereby settling melanoma on EI monolayer (Fig. 8F). These results indicate that thrombin-mediated Fg conversion to Fn plays influential roles in melanoma–PMN–EC adhesion, mechanistically via the receptors of ICAM-1 and  $\alpha_v\beta_3$  on melanoma.

In summary, our studies showed that fibrin(ogen) could enhance the binding between PMNs and melanoma cells under shear conditions; ICAM-1 and  $\alpha_v\beta_3$  contributed to this process. Under hydrodynamic conditions, ICAM-1/Fn/Mac-1 and ICAM-1/LFA-1 bonds promoted effective cell–cell tethering and prolonged initial cell–cell contact duration, whereas  $\alpha_v\beta_3$ /Fn/Mac-1 bonds enhanced cell–cell sustained adhesion, demonstrating cooperative roles of ICAM-1 and  $\alpha_v\beta_3$  in tumor cell–PMN adhesion in the presence of Fn (Fig. 9).

## Discussion

In the current study, we have demonstrated that: 1) melanoma cells and PMNs can attach to immobilized Fg and Fn under venous flow conditions.  $\alpha_v\beta_3$  on Lu1205 melanoma cells and Mac-1 on PMNs are the primary receptors for binding to immobilized Fg and Fn; 2)




**FIGURE 7.** Receptors on Lu1205 cells other than ICAM-1 and  $\alpha_v\beta_3$  played marginal roles in Lu1205 and PMN heterotypic aggregation. *A*, The effect of ICAM-1 and CD51/61 blocking on Lu1205 and PMN heterotypic aggregation in the presence of Fn. Fibrin solution was made from 0.053 U/ml thrombin, 1.5 mg/ml Fg, and 4 mM GPRP. Values are means  $\pm$  SEM from three independent experiments. *B*, The effect of Mac-1 blocking on Lu1205 and PMN heterotypic aggregation in the presence or absence of Fg. Values are means  $\pm$  SEM ( $n > 3$ ). *C*, The abilities of melanoma cells with different metastatic potentials and different receptor expressions to bind to PMNs in the presence of Fn. Lu1205 expresses high levels of ICAM-1 and  $\alpha_v\beta_3$ , A375m expresses median levels of ICAM-1 and low levels of  $\alpha_v\beta_3$ , and WM35 expresses median levels of  $\alpha_v\beta_3$  and low levels of ICAM-1. Cells were pretreated with 1 U/ml chondroitinase AC II for 60 min at 37°C to remove terminal chondroitin sulfate and dermatan sulfate residues (on CD44). Values are means  $\pm$  SEM ( $n > 3$ ).

soluble Fg and Fn can enhance the heteroaggregation between melanoma cells and PMNs as the bridging molecules, although the kinetics of Fn-mediated binding is different from that of Fg; and 3)  $\alpha_v\beta_3$  and ICAM-1 cooperate to mediate PMN and melanoma cell binding in the presence of soluble Fg or Fn under flow conditions, where ICAM-1 mediates initial tethering, whereas  $\alpha_v\beta_3$  contributes to firm adhesion (Fig. 9).

*Soluble fibrin or fibrinogen, as a dimeric molecule, supports both initial capture and stable adhesion of melanoma cells to neutrophils under flow conditions*

Tumor cells and PMNs in a cone-plate viscometer are subject to a uniform shear field, resulting in mutual collisions (9, 17). This millisecond scale of shear-dependent encounter has been shown to be able to mediate receptor–ligand bond formation between melanoma cells and PMNs in the absence of fibrin(ogen) (16). Increasing shear force opposes bond formation by decreasing the cell contact duration but by increasing the cell–cell contact area. If the ensemble strength of receptor–ligand bonds outweighs tensile force, the heterotypic tumor cell–PMN aggregates form; otherwise, the aggregates dissociate. We now have shown that Fg mainly enhances the initial tethering of melanoma cells to PMNs, whereas Fn profoundly changes the kinetics of cell–cell binding, stabilizing melanoma–PMN doublets for several minutes. Results from Fig. 5A show that the extent of aggregation was increased with an increasing amount of Fg. However, this trend would be reversed after the concentration of soluble Fg reaches a certain level, because high amounts of free ligands tend to saturate counterreceptors on mutual cell types instead of bridging two cell receptors.


Tumor cell-mediated activation of coagulation cascade is a key factor for hematogenous metastasis. In this process, Fn plays various roles. Fn has also been shown to support migration of different cell types, including monocytes and fibroblasts. Its proteolytic products are chemotactic and angiogenic reagents. In addition, Fn is related to tumor stroma formation and invasion. Because thrombin generation is a determinant for Fn formation, it is interesting to see how thrombin affects tumor cell adhesion. ELISA results have shown that thrombin in plasma was increased when Lu1205 and A2058 melanoma cells were cultured or cocultured with HUVECs (data not shown). Thrombin enhances tumor cell adhesion to platelets and facilitated tumor cell metastasis in vivo (33). Thrombin may also promote tumor cell metastasis through an Fg-independent mechanism (7). The possible effect of thrombin on Fg can be described using a Michaelis-Menten scheme in which Fg binds to thrombin with certain rates and affinities (34). The rate of conversion of Fg to Fn is dependent on the concentration of thrombin. To understand the potential role of thrombin on conversion of Fg to Fn, we separated Fn-digested products with SDS-PAGE. Most of  $\alpha$ - and  $\beta$ -chains were cleaved by thrombin to produce lower m.w. products. This demonstrated the functionality of thrombin and purity of Fn produced by 2 U/ml thrombin. Thrombin concentration of 2 U/ml was assumed to convert all the 1.5 mg/ml Fg to Fn within a 10-min incubation time period (8). To probe the sensitivity of cell adhesion to Fn concentration, we have varied the concentrations of thrombin for Fn generation. We tried a range of thrombin concentrations, varying from 0.0265 U/ml to 2 U/ml. Results have shown that Fn solution made from 0.053 U/ml thrombin significantly enhanced  $\alpha_v\beta_3$ -mediated firm adhesion between melanoma cells and PMNs, suggesting that cryptic  $\alpha_v\beta_3$  binding sites are exposed by thrombin catalysis. When thrombin level was further increased to 2 U/ml from 0.053 U/ml, only the initial tethering phase was affected. This could imply that more ICAM-1 binding sites were available when melanoma cells were subject to higher concentration of thrombin. Therefore, we speculate that  $\alpha_v\beta_3$  binding sites could be first exposed upon conversion of Fg to Fn, whereas it may require higher dose of thrombin to make ICAM-1 binding sites accessible. This hypothesis needs further biochemical analysis for proof. The fact that Fn has a higher affinity than Fg for receptors on melanoma cells agrees with the results from other cell types in which CD44 on carcinoma cells had a higher affinity for Fn than Fg (8, 9, 31). Thrombin may have



**FIGURE 8.** Relative contributions of  $\alpha_v\beta_3$  and ICAM-1 to fibrin(ogen)-mediated PMN–melanoma–EC interaction under flow in parallel plate. *A–D*, The number of PMNs contacting with Lu1205 or WM35 cells for >1 s (initial tethering), 3 s (transient adhesion), and 10 s (long-term adhesion): fibrin(ogen) enhanced PMN and melanoma cell interaction (*A*). Data are means  $\pm$  SEM ( $n > 3$ ). \* $p < 0.05$  compared with control; § $p < 0.05$  compared with fibrinogen cases. Roles of  $\alpha_v\beta_3$  and ICAM-1 in different phases of adhesion of PMNs to Lu1205 cells under shear rate of 62.5 s<sup>-1</sup> (*B*) or 20 s<sup>-1</sup> (*C*). Values are means  $\pm$  SEM ( $n > 3$ ). \* $p < 0.05$  compared with fibrin only cases. *D*, PMNs bind weakly to nonmetastatic melanoma cell line WM35. Values are means  $\pm$  SEM ( $n > 3$ ). \* $p < 0.05$  compared with Lu1205 cases. The adhesion efficiency of Lu1205 cells binding to EI cells for >1 s (initial tethering), 3 s (transient adhesion), and 5 s (long-term adhesion) as a result of Lu1205 and PMN collisions: role of soluble fibrin(ogen) on melanoma adhesion efficiency to EI cells via PMNs at shear rate of 62.5 s<sup>-1</sup> (*E*); roles of  $\alpha_v\beta_3$  and ICAM-1 in Lu1205 adhesion at the shear rate of 62.5 s<sup>-1</sup> (*F*). \* $p < 0.05$  compared with the Lu1205+Fn case. Values are mean  $\pm$  SEM ( $n > 3$ ).

a potential effect on  $\beta_2$  integrin expression on PMNs. Therefore, the residual thrombin in our Fn solution may itself affect PMN–melanoma cell aggregation. However, it should be noted that all PMNs used in the current studies were prestimulated with fMLF, which boosted the Mac-1 expression to maximal levels when we conducted cone-plate and parallel plate assays (35). In this way, we can preclude the possibility that the observed Fn-mediated aggregation was the pseudoimage of thrombin-stimulated PMN–melanoma binding.

The concept of Fn-mediated melanoma cell and PMN aggregation under flow conditions is intriguing. Several earlier studies demonstrated that Fn might mediate platelet–tumor cell adhesion (3, 5). These platelet–tumor cell microemboli may tether tumor cells to the endothelial wall, thereby contributing to tumor metastasis. Other similar evidence of Fn-mediated cell–cell interaction came from the studies of leukocyte adhesion to EC, where Mac-1 on PMNs and ICAM-1 on the ECs may be the respective receptors for Fg (6, 25). To our best knowledge, the current study



**FIGURE 9.** A model concluded from the present studies on PMN–Lu1205 heterotypic aggregation in the presence of Fn under shear conditions. Fn-bound ICAM-1 quickly binds to Mac-1 on PMNs, thereby increasing the cell–cell contact duration and allowing Fn-bound  $\alpha_v\beta_3$  to mediate firm adhesion between PMN–Lu1205.

has first shown that thrombin-regulated Fn formation can enhance PMN and melanoma cell aggregation under flow conditions. An earlier study reported that soluble Fn would actually reduce the monocyte adherence to A375 melanoma cells under flow conditions (22). However, there are several differences between these studies and the current work. The melanoma cell line A375 used in Biggerstaff et al. (22) was reported to express ICAM-1 but not  $\alpha_v\beta_3$ , whereas Lu1205 cells used in the current study express  $\alpha_v\beta_3$  in addition to ICAM-1, which was shown to have a high affinity for fibrin(ogen)-mediated firm adhesion of melanoma cells. In addition, experimental methods used in Biggerstaff et al. (22) were different from the current study. In Biggerstaff et al. (22), monocytes were perfused over tumor cell monolayer for 1 h in a parallel-plate flow chamber. In contrast, we quantified PMN–Lu1205 heterotypic aggregation within short time periods (several minutes). In a cone-plate assay, both Fn and Fg were able to significantly enhance aggregation between  $\alpha_v\beta_3$ -blocked Lu1205 cells and PMNs within 5 min shear (Fig. 6A–C). By 60 s, Fn and Fg increased percentage of aggregation from 45 to 60%; by 5 min, Fn and Fg increased percentage of aggregation from 27 to 37 and 35%, respectively. This result implies that Fn promotes Mac-1/ICAM-1 bond formation by possibly adding extra binding sites between Mac-1 and ICAM-1. However, during the long-term shear, Mac-1/Fn/ICAM-1 bonds dissociate (22). Without  $\alpha_v\beta_3$ , Fn-occupied ICAM-1 or Mac-1 could become obstacles to normal Mac-1/ICAM-1 bond formation, thereby reducing aggregation.

#### *$\alpha_v\beta_3$ integrin contributes to fibrin-mediated prolonged stable adhesion between melanoma cells and PMNs*

Melanoma cells do not express selectins or sLe<sup>x</sup> sugar groups at the levels necessary for cells to attach to the endothelial wall under venous flow conditions. Recent studies have shown that fibrinolytic factors and Fn deposition were both associated with hematogenous metastasis of murine melanoma cells (7). Fn is likely to be deposited on the surface of endothelium or platelets upon inflammation or tissue damage. Melanoma cells attach to Fg via  $\alpha_v\beta_3$  and  $\alpha_v\beta_1$ , to von Willebrand Factor via  $\alpha_v\beta_3$  and to fibronectin via  $\alpha_v\beta_3$ ,  $\alpha_v\beta_1$  and  $\alpha_5\beta_1$ , respectively, under static conditions. In addition, the current study using a parallel-plate flow chamber to examine Lu1205 melanoma cells binding to immobilized fibrin(ogen) has indicated that  $\alpha_v\beta_3$  was a major contributing receptor for Fg and Fn binding especially under low shear rates, which agrees well with a previous study on M21 melanoma cells adhesion to fibrin(ogen) (11). More importantly, melanoma cells could engage with fibrin(ogen) firmly without apparent cell rolling, which is

potentially important for the arrest of sLe<sup>x</sup>-negative melanoma cells the endothelial cells under venous flow conditions.

The expression of  $\alpha_v\beta_3$  is associated with malignant phenotype of tumor, which promotes tumor cells, ECs, and fibroblast migration and invasion by interacting with fibrin(ogen) and its plasminogen-lytic products (36, 37). When  $\alpha_v\beta_3$  on melanoma cells were blocked, Fn-mediated sustained aggregation between melanoma cells and PMNs was almost obligated. This demonstrated that  $\alpha_v\beta_3$  have a high affinity for Fn-mediated stable-firm adhesion between PMNs and melanoma cells under hydrodynamic conditions. The high affinity of  $\alpha_v\beta_3$  for fibrin(ogen) is evident from biochemical and structural analysis. There are three putative  $\alpha_v\beta_3$  binding sites on Fg (13), which are RGDS at the C terminus of  $\alpha$ -chain A $\alpha$  572–575, RGDF at N terminus of  $\alpha$ -chain A $\alpha$  95–98, and dodecapeptide at the C terminus of  $\gamma$ -chain  $\gamma$ 400–411. These domains bind to immobilized  $\alpha_v\beta_3$  so strongly that they do not dissociate once they bind. In particular, RGDS site at A $\alpha$  572–575 has stronger affinity for  $\alpha_v\beta_3$  than dodecapeptide. Thrombin treatment of Fg on its  $\alpha$ -chain may induce conformational change and expose these RGD sites that are inaccessible to integrins in native structures. In addition, fibrin(ogen) may activate  $\alpha_v\beta_3$  and induce cluster of integrins on cell–cell contact regions, further increasing the stability of cell aggregates (38). In agreement with our heterotypic aggregation studies and model about fibrin(ogen)-mediated PMN-dependent melanoma adhesion, soluble Fg enhanced the melanoma cell arrest to immobilized Fg, serving as a cross-linking ligand for  $\alpha_v\beta_3$  between attached and circulating tumor cells (11). More in-depth kinetic analysis of  $\alpha_v\beta_3$ –fibrin(ogen) interaction is needed to better understand the bond strength.

#### *Mac-1 on PMNs serves as a counterreceptor for ICAM-1 and fibrin(ogen)*

Mac-1 on leukocytes, especially PMNs, is a high affinity receptor for fibrin(ogen), mediating PMN adhesion to inflamed ECs (39). Previously, Mac-1 was shown to mediate PMN homotypic aggregation under venous flow conditions (40). The current study demonstrates that Mac-1 also played roles in Fn-mediated heterotypic aggregation between melanoma cells and PMNs. Fg contains many recognition sites for Mac-1 within D domain, including two peptide sequences,  $\gamma$ 190–202 and  $\gamma$ 377–395 (41), which are called P1 and P2. Mac-1 may noncovalently bind to sites other than P1 and P2 (42). However, Mac-1 has stronger avidity for soluble Fn or immobilized Fg. This was suggested by the cryptic theory where pulling out P2 region from the central domain of Fg  $\gamma$ -module would regulate the binding affinity of Fg for Mac-1. This partially explains our melanoma–PMN aggregation results showing that Fn mediates stronger binding than Fg does. Although some studies also showed that PMNs have a higher possibility to form long-lasting bonds with immobilized ICAM-1 than immobilized Fg (27), we cannot rule out the possibility that Fn could enhance PMNs and tumor cell binding by inserting additional bonds between Mac-1 and ICAM-1 instead of replacing the existing bonds.

Collectively, our study demonstrated that PMN could tether melanoma cells to the vascular endothelial wall through the formation of shear-resist bonds. The most intriguing part of the current study is to demonstrate how a tumor microenvironment affects immune cell responses and cancer cell adhesion, especially when being subject to thrombin-regulated fibrin(ogen) immunoeediting. This is one of several possible settings in which tumor cells can capitalize on hemostatic components for the metastasis. This study has shed light on possible molecular mechanisms in fibrin(ogen)-

mediated PMN–melanoma interactions under flow conditions and potential therapeutic importance of Fn receptors in melanoma cell adhesion.

*Potential roles of selectin ligands, P-selectin, L-selectin, and VCAM-1 on melanoma cell adhesion*

Conventionally, the binding of P-selectin to leukocytes requires sialylated and fucosylated carbohydrates on PSGL-1. This binding is supported by sLe<sup>x</sup>-like motifs on O-linked glycans of PSGL-1. Analysis of human metastatic melanoma lesions showed an elevated expression of sLe<sup>a</sup> and sLe<sup>x</sup> (43). However, further exploration of the interaction between melanoma and ECs revealed that melanoma may use other proteoglycan and oligosaccharide to bind to P-selectin and E-selectin, including heparin sulfate under flow (44). Using mAbs failed to detect any apparent expression of sLe<sup>x</sup> on small cell lung cancer and neuroblastoma cells (45). Therefore, it is of interest to compare PMN-mediated, Fn-mediated, and selectin-mediated melanoma adhesion to ECs under flow. From our adhesion studies using parallel-plate assays, most melanoma arresting onto the EI monolayer were associated with PMNs. In addition, Fn enhanced PMN-facilitated melanoma adhesion in higher magnitude than did melanoma direct adhesion to ECs (data not shown). Despite importance of PMN and Fn, selectins and VCAM-1 on ECs should not be ignored. VLA-4, which is expressed by melanoma, has ligand VCAM-1 on ECs. Upregulation of VCAM-1 expression was associated with increased experimental metastasis in vivo (46, 47). VCAM-1/VLA-4 interaction-mediated melanoma adhesion to ECs under low shear condition (48, 49). Surprisingly, a high level of VLA-4 may restrict cells to detach as a prelude to invasion, thereby inhibiting tumor cell metastasis. Selectin roles in melanoma metastasis were also demonstrated in animal models. For example, L-selectin plays a role in nonlymphoma tumor cell localization to lymph node (50). Although P-selectin was traditionally considered as a receptor for platelet–melanoma binding, its role in EC–melanoma binding was studied in P-selectin-deficient mice in which lung metastasis was significantly attenuated (51).

Tether behavior exhibited by fibrin(ogen)–ICAM-1 bonds may be an indication that in absence of selectin ligands, melanoma cells may explore other mechanisms for their successful transmigration. There have been no biomechanical models or biochemical basis for this interaction so far. Fast association and dissociation rates are prerequisites for PSGL-1–P-selectin-mediated PMN tether to ECs under flow conditions. Such a rapid kinetic property was not witnessed in Mac-1/Fn and  $\alpha_v\beta_3$ /Fn binding measured by surface plasmon resonance. Based on the current study, it is may be possible that effective tumor adhesion to PMN and ECs in the presence of multivalent plasma protein does not require a rapid association rate. It is intriguing to think whether it is unique among Fn-mediated cell–cell interactions or in a similar in magnitude to reported selectin–PSGL-1 binding.

*Potential roles of receptors in melanoma metastasis*

During their passage through the EC wall, tumor cells undergo extensive interactions with host cells including PMNs. PMNs cause a dense Fn accumulation around them under flow conditions with a platelet-independent mechanism in which Mac-1 may play a role (52). This may contribute to binding of melanoma to PMNs and arrest of melanoma to ECs. A dose- and time-dependent increase in ICAM-1 on melanoma by TNF- $\alpha$  treatment resulted in enhancement of in vivo metastasis (53). This TNF- $\alpha$  inducible increase in metastasis could be reversed by knocking down ICAM-1 on cell surface. Apparent contradictory results came from in vivo studies with ICAM-1 knockout mice, which showed that ICAM-1

deletion enhanced pulmonary metastasis (54, 55). However, this genetic deletion was conducted on the acceptor mice side but not on the melanoma side. Because T cells use receptors for ICAM-1 to bind/migrate into tissue and mediate tumor killing processes, reduced ICAM-1 expression may inhibit T cell migration and T cell-mediated tumor rejection. Most likely, the reduced tumor killing might compensate for the reduced tumor adhesion to ECs in ICAM-1 knockout mice, leading to the enhanced melanoma lesions observed in tissue outside the vasculature. A better understanding of involvement of receptors in melanoma adhesion to ECs via Fn and PMNs in circulation may come from examination of melanoma arrest within vasculature.

## Acknowledgments

We thank Dr. M. Herlyn (Wistar Institute, Philadelphia, PA) for providing WM35 melanoma cells and Dr. Scott Simon (University of California Davis, Davis, CA) for providing EI cells. We also thank Dr. Hari Muddana and Dr. Shile Liang for many helpful discussions and technical support.

## Disclosures

The authors have no financial conflicts of interest.

## References

- Liang, S., M. J. Slattery, and C. Dong. 2005. Shear stress and shear rate differentially affect the multi-step process of leukocyte-facilitated melanoma adhesion. *Exp. Cell Res.* 310: 282–292.
- Biggerstaff, J. P., N. Seth, A. Amirhosravi, M. Amaya, S. Fogarty, T. V. Meyer, F. Siddiqui, and J. L. Francis. 1999. Soluble fibrinogen augments platelet/tumor cell adherence in vitro and in vivo, and enhances experimental metastasis. *Clin. Exp. Metastasis* 17: 723–730.
- Felding-Habermann, B., R. Habermann, E. Saldívar, and Z. M. Ruggeri. 1996. Role of beta3 integrins in melanoma cell adhesion to activated platelets under flow. *J. Biol. Chem.* 271: 5892–5900.
- Mueller, B. M., R. A. Reisfeld, T. S. Edgington, and W. Ruf. 1992. Expression of tissue factor by melanoma cells promotes efficient hematogenous metastasis. *Proc. Natl. Acad. Sci. USA* 89: 11832–11836.
- Biggerstaff, J. P., N. B. Seth, T. V. Meyer, A. Amirhosravi, and J. L. Francis. 1998. Fibrin monomer increases platelet adherence to tumor cells in a flowing system: a possible role in metastasis? *Thromb. Res.* 92(6, Suppl 2): S53–S58.
- Languino, L. R., J. Plescia, A. Duperray, A. A. Brian, E. F. Plow, J. E. Geltosky, and D. C. Altieri. 1993. Fibrinogen mediates leukocyte adhesion to vascular endothelium through an ICAM-1-dependent pathway. *Cell* 73: 1423–1434.
- Palumbo, J. S., K. W. Kombrinck, A. F. Drew, T. S. Grimes, J. H. Kiser, J. L. Degen, and T. H. Bugge. 2000. Fibrinogen is an important determinant of the metastatic potential of circulating tumor cells. *Blood* 96: 3302–3309.
- Alves, C. S., M. M. Burdick, S. N. Thomas, P. Pawar, and K. Konstantopoulos. 2008. The dual role of CD44 as a functional P-selectin ligand and fibrin receptor in colon carcinoma cell adhesion. *Am. J. Physiol. Cell Physiol.* 294: C907–C916.
- Goldsmith, H. L., T. A. Quinn, G. Drury, C. Spanos, F. A. McIntosh, and S. I. Simon. 2001. Dynamics of neutrophil aggregation in couette flow revealed by videomicroscopy: effect of shear rate on two-body collision efficiency and doublet lifetime. *Biophys. J.* 81: 2020–2034.
- Liang, S., M. J. Slattery, D. Wagner, S. I. Simon, and C. Dong. 2008. Hydrodynamic shear rate regulates melanoma-leukocyte aggregation, melanoma adhesion to the endothelium, and subsequent extravasation. *Ann. Biomed. Eng.* 36: 661–671.
- Pilch, J., R. Habermann, and B. Felding-Habermann. 2002. Unique ability of integrin alpha(v)beta 3 to support tumor cell arrest under dynamic flow conditions. *J. Biol. Chem.* 277: 21930–21938.
- Felding-Habermann, B., Z. M. Ruggeri, D. A. Cheresch, and D. A. Cheresch. 1992. Distinct biological consequences of integrin alpha v beta 3-mediated melanoma cell adhesion to fibrinogen and its plasmic fragments. *J. Biol. Chem.* 267: 5070–5077.
- Yokoyama, K., H. P. Erickson, Y. Ikeda, and Y. Takada. 2000. Identification of amino acid sequences in fibrinogen gamma-chain and tenascin C C-terminal domains critical for binding to integrin alpha v beta 3. *J. Biol. Chem.* 275: 16891–16898.
- Altieri, D. C., A. Duperray, J. Plescia, G. B. Thornton, and L. R. Languino. 1995. Structural recognition of a novel fibrinogen  $\gamma$  chain sequence (117–133) by intercellular adhesion molecule-1 mediates leukocyte-endothelium interaction. *J. Biol. Chem.* 270: 696–699.
- Lishko, V. K., N. P. Podolnikova, V. P. Yakubenko, S. Yakovlev, L. Medved, S. P. Yadav, and T. P. Ugarova. 2004. Multiple binding sites in fibrinogen for integrin alphaMbeta2 (Mac-1). *J. Biol. Chem.* 279: 44897–44906.
- Liang, S., C. Fu, D. Wagner, H. Guo, D. Zhan, C. Dong, and M. Long. 2008. Two-dimensional kinetics of  $\beta_2$ -integrin and ICAM-1 bindings between neutrophils and melanoma cells in a shear flow. *Am. J. Physiol. Cell Physiol.* 294: C743–C753.

17. Hentzen, E. R., S. Neelamegham, G. S. Kansas, J. A. Benanti, L. V. McIntire, C. W. Smith, and S. I. Simon. 2000. Sequential binding of CD11a/CD18 and CD11b/CD18 defines neutrophil capture and stable adhesion to intercellular adhesion molecule-1. *Blood* 95: 911–920.
18. Liang, S., A. Sharma, H. H. Peng, G. Robertson, and C. Dong. 2007. Targeting mutant (V600E) B-Raf in melanoma interrupts immunoeediting of leukocyte functions and melanoma extravasation. *Cancer Res.* 67: 5814–5820.
19. Alves, C. S., S. Yakovlev, L. Medved, and K. Konstantopoulos. 2009. Bio-molecular characterization of CD44-fibrin(ogen) binding: distinct molecular requirements mediate binding of standard and variant isoforms of CD44 to immobilized fibrin(ogen). *J. Biol. Chem.* 284: 1177–1189.
20. Simon, S. I., Y. Hu, D. Vestweber, and C. W. Smith. 2000. Neutrophil tethering on E-selectin activates beta 2 integrin binding to ICAM-1 through a mitogen-activated protein kinase signal transduction pathway. *J. Immunol.* 164: 4348–4358.
21. Gopalan, P. K., C. W. Smith, H. Lu, E. Berg, L. V. McIntire, and S. I. Simon. 1997. Neutrophil CD18-dependent arrest on intercellular adhesion molecule-1 (ICAM-1) in shear flow can be activated through L-selectin. *J. Immunol.* 158: 367–375.
22. Biggerstaff, J. P., B. Weidow, J. Vidosh, J. Dexheimer, S. Patel, and P. Patel. 2006. Soluble fibrin inhibits monocyte adherence and cytotoxicity against tumor cells: implications for cancer metastasis. *Thromb. J.* 4: 12.
23. Hoskins, M. H., and C. Dong. 2006. Kinetics analysis of binding between melanoma cells and neutrophils. *Mol. Cell. Biomech.* 3: 79–87.
24. Jadhav, S., B. S. Bochner, and K. Konstantopoulos. 2001. Hydrodynamic shear regulates the kinetics and receptor specificity of polymorphonuclear leukocyte-cytocollon carcinoma cell adhesive interactions. *J. Immunol.* 167: 5986–5993.
25. Duperray, A., L. R. Languino, J. Plescia, A. McDowall, N. Hogg, A. G. Craig, A. R. Berendt, and D. C. Altieri. 1997. Molecular identification of a novel fibrinogen binding site on the first domain of ICAM-1 regulating leukocyte-endothelium bridging. *J. Biol. Chem.* 272: 435–441.
26. Kuijper, P. H., H. I. Gallardo Torres, J. A. van der Linden, J. W. Lammers, J. J. Sixma, J. J. Zwaginga, and L. Koenderman. 1997. Neutrophil adhesion to fibrinogen and fibrin under flow conditions is diminished by activation and L-selectin shedding. *Blood* 89: 2131–2138.
27. Kadash, K. E., M. B. Lawrence, and S. L. Diamond. 2004. Neutrophil string formation: hydrodynamic thresholding and cellular deformation during cell collisions. *Biophys. J.* 86: 4030–4039.
28. Li, P., P. Selvaraj, and C. Zhu. 1999. Analysis of competition binding between soluble and membrane-bound ligands for cell surface receptors. *Biophys. J.* 77: 3394–3406.
29. Yakovlev, S., S. Gorlatov, K. Ingham, and L. Medved. 2003. Interaction of fibrin (ogen) with heparin: further characterization and localization of the heparin-binding site. *Biochemistry* 42: 7709–7716.
30. Neelamegham, S., A. D. Taylor, J. D. Hellums, M. Dembo, C. W. Smith, and S. I. Simon. 1997. Modeling the reversible kinetics of neutrophil aggregation under hydrodynamic shear. *Biophys. J.* 72: 1527–1540.
31. Konstantopoulos, K., and S. N. Thomas. 2009. Cancer cells in transit: the vascular interactions of tumor cells. *Annu. Rev. Biomed. Eng.* 11: 177–202.
32. Slattery, M. J., S. Liang, and C. Dong. 2005. Distinct role of hydrodynamic shear in leukocyte-facilitated tumor cell extravasation. *Am. J. Physiol. Cell Physiol.* 288: C831–C839.
33. Nierodzick, M. L., F. Kajumo, and S. Karpatkin. 1992. Effect of thrombin treatment of tumor cells on adhesion of tumor cells to platelets in vitro and tumor metastasis in vivo. *Cancer Res.* 52: 3267–3272.
34. Scheraga, H. A. 2004. The thrombin-fibrinogen interaction. *Biophys. Chem.* 112: 117–130.
35. Neelamegham, S., A. D. Taylor, A. R. Burns, C. W. Smith, and S. I. Simon. 1998. Hydrodynamic shear shows distinct roles for LFA-1 and Mac-1 in neutrophil adhesion to intercellular adhesion molecule-1. *Blood* 92: 1626–1638.
36. Gailit, J., C. Clarke, D. Newman, M. G. Tonnesen, M. W. Mosesson, and R. A. F. Clark. 1997. Human fibroblasts bind directly to fibrinogen at RGD sites through integrin alpha(v)beta3. *Exp. Cell Res.* 232: 118–126.
37. Leavesley, D. I., G. D. Ferguson, E. A. Wayner, and D. A. Cheresh. 1992. Requirement of the integrin beta 3 subunit for carcinoma cell spreading or migration on vitronectin and fibrinogen. *J. Cell Biol.* 117: 1101–1107.
38. Cluzel, C., F. Saltel, J. Lussi, F. Paulhe, B. A. Imhof, and B. Wehrle-Haller. 2005. The mechanisms and dynamics of (alpha)v(beta)3 integrin clustering in living cells. *J. Cell Biol.* 171: 383–392.
39. Diamond, M. S., and T. A. Springer. 1993. A subpopulation of Mac-1 (CD11b/CD18) molecules mediates neutrophil adhesion to ICAM-1 and fibrinogen. *J. Cell Biol.* 120: 545–556.
40. Simon, S. I., J. D. Chambers, E. Butcher, and L. A. Sklar. 1992. Neutrophil aggregation is beta 2-integrin- and L-selectin-dependent in blood and isolated cells. *J. Immunol.* 149: 2765–2771.
41. Mosesson, M. W. 2005. Fibrinogen and fibrin structure and functions. *J. Thromb. Haemost.* 3: 1894–1904.
42. Yakovlev, S., L. Zhang, T. Ugarova, and L. Medved. 2005. Interaction of fibrin (ogen) with leukocyte receptor alpha M beta 2 (Mac-1): further characterization and identification of a novel binding region within the central domain of the fibrinogen gamma-module. *Biochemistry* 44: 617–626.
43. Kageshita, T., S. Hirai, T. Kimura, N. Hanai, S. Ohta, and T. Ono. 1995. Association between sialyl Lewis(x) expression and tumor progression in melanoma. *Cancer Res.* 55: 1748–1751.
44. Ma, Y. Q., and J. G. Geng. 2000. Heparan sulfate-like proteoglycans mediate adhesion of human malignant melanoma A375 cells to P-selectin under flow. *J. Immunol.* 165: 558–565.
45. Stone, J. P., and D. D. Wagner. 1993. P-selectin mediates adhesion of platelets to neuroblastoma and small cell lung cancer. *J. Clin. Invest.* 92: 804–813.
46. Okahara, H., H. Yagita, K. Miyake, and K. Okumura. 1994. Involvement of very late activation antigen 4 (VLA-4) and vascular cell adhesion molecule 1 (VCAM-1) in tumor necrosis factor alpha enhancement of experimental metastasis. *Cancer Res.* 54: 3233–3236.
47. Vidal-Vanaclocha, F., G. Fantuzzi, L. Mendoza, A. M. Fuentes, M. J. Anasagasti, J. Martín, T. Carrascal, P. Walsh, L. L. Reznikov, S. H. Kim, et al. 2000. IL-18 regulates IL-1b-dependent hepatic melanoma metastasis via vascular cell adhesion molecule-1. *Proc. Natl. Acad. Sci. USA.* 97: 734–739.
48. Liang, S., and C. Dong. 2008. Integrin VLA-4 enhances sialyl-Lewis(x)/a-negative melanoma adhesion to and extravasation through the endothelium under low flow conditions. *Am. J. Physiol. Cell Physiol.* 295: C701–C707.
49. Mould, A. P., J. A. Askari, S. E. Craig, A. N. Garratt, J. Clements, and M. J. Humphries. 1994. Integrin  $\alpha_4\beta_1$ -mediated melanoma cell adhesion and migration on vascular cell adhesion molecule-1 (VCAM-1) and the alternatively spliced IIIc3 region of fibronectin. *J. Biol. Chem.* 269: 27224–27230.
50. Qian, F., D. Hanahan, and I. L. Weissman. 2001. L-selectin can facilitate metastasis to lymph nodes in a transgenic mouse model of carcinogenesis. *Proc. Natl. Acad. Sci. USA* 98: 3976–3981.
51. Ludwig, R. J., B. Boehme, M. Podda, R. Henschler, E. Jager, C. Tandl, W. H. Boehncke, T. M. Zollner, R. Kaufmann, and J. Gille. 2004. Endothelial P-selectin as a target of heparin action in experimental melanoma lung metastasis. *Cancer Res.* 64: 2743–2750.
52. Goel, M. S., and S. L. Diamond. 2001. Neutrophil enhancement of fibrin deposition under flow through platelet-dependent and -independent mechanisms. *Arterioscler. Thromb. Vasc. Biol.* 21: 2093–2098.
53. Miele, M. E., C. F. Bennett, B. E. Miller, and D. R. Welch. 1994. Enhanced metastatic ability of TNF-alpha-treated malignant melanoma cells is reduced by intercellular adhesion molecule-1 (ICAM-1, CD54) antisense oligonucleotides. *Exp. Cell Res.* 214: 231–241.
54. Yamada, M., K. Yanaba, M. Hasegawa, Y. Matsushita, M. Horikawa, K. Komura, T. Matsushita, A. Kawasuji, T. Fujita, K. Takehara, et al. 2006. Regulation of local and metastatic host-mediated anti-tumour mechanisms by L-selectin and intercellular adhesion molecule-1. *Clin. Exp. Immunol.* 143: 216–227.
55. Marvin, M. R., J. C. Southall, S. Trokhan, C. DeRosa, and J. Chabot. 1998. Liver metastases are enhanced in homozygous deletionally mutant ICAM-1 or LFA-1 mice. *J. Surg. Res.* 80: 143–148.